

PRV

PATENT- OCH REGISTRERINGSVERKET
Patentavdelningen

PCT/SE 03 / 0.1078

#2

Intyg Certificate

Härmed intygas att bifogade kopior överensstämmer med de handlingar som ursprungligen ingivits till Patent- och registreringsverket i nedannämnda ansökan.

Ansökan ingavs ursprungligen på engelska.


This is to certify that the annexed is a true copy of the documents as originally filed with the Patent- and Registration Office in connection with the following patent application.

The application was originally filed in English.

(71) Sökande AstraZeneca AB, Södertälje SE
Applicant (s)

(21) Patentansökningsnummer 0201976-8
Patent application number

(86) Ingivningsdatum 2002-06-24
Date of filing



Stockholm, 2003-07-01

För Patent- och registreringsverket
For the Patent- and Registration Office

Sonia André

Avgift
Fee

PRIORITY DOCUMENT
SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH
RULE 17.1(a) OR (b)

**PATENT- OCH
REGISTRERINGSVERKET
SWEDEN**

Postadress/Adress
Box 5055
S-102 42 STOCKHOLM

Telefon/Phone
+46 8 782 25 00
Vx 08-782 25 00

Telex
17978
PATOREG S

Telefax
+46 8 666 02 86
08-666 02 86

NOVEL COMPOUNDS


The present invention relates to compounds and compositions for treating diseases associated with cysteine protease activity. The compounds are reversible inhibitors of cysteine proteases S, K, F, L and B. Of particular interest are diseases associated with Cathepsin S. In addition this invention also discloses processes for the preparation of such inhibitors.

BACKGROUND OF THE INVENTION

Cathepsin S is a member of the papain superfamily of cysteine proteases which also encompasses Cathepsins B, H, L, O and K. Cathepsin S plays a key role in the processing of invariant chain in MHC class II complexes allowing the complex to associate with antigenic peptides. MHC class II complexes are then transported to the surface of the cell for presentation to effector cells such as T cells. The process of antigen presentation is a fundamental step in initiation of the immune response. In this respect inhibitors of cathepsin S could be useful agents in the treatment of inflammation and immune disorders such as, but not limited to, asthma, rheumatoid arthritis, multiple sclerosis and Crohn's disease. Cathepsin S has also been implicated in a variety of other diseases involving extracellular proteolysis such as the development of emphysema in COPD through degradation of elastin and in Alzheimers disease.

Other Cathepsins notably K and L have been shown to degrade bone collagen and other bone matrix proteins. Inhibitors of these cysteine proteases would be expected to be useful in the treatment of diseases involving bone resorption such as osteoporosis.

The present invention therefore provides a compound of formula (I)



(I)

in which:

X is N or CA where A is hydrogen, halogen, CHR^2R^3 , OR^2 , NR^2R^3 , SR^2 ;

5 R^2 and R^3 are independently hydrogen, C_{1-6} alkyl or C_{3-6} cycloalkyl both of which can optionally contain one or more O, S or NR^4 groups where R^4 is hydrogen or C_{1-6} alkyl, and can be optionally substituted by aryl, heteroaryl, NR^5R^6 where R^5 and R^6 together with the nitrogen atom to which they are attached form a 4-7 membered ring optionally containing a
10 further O, S, NR^4 , or R^2 and R^3 together with the nitrogen atom to which they are attached form a 4-7 membered ring optionally containing a further O, S, NR^4 group, or R^2 and R^3 are aryl or heteroaryl groups, both aryl and heteroaryl groups being optionally substituted by halogen, amino, hydroxy, cyano, nitro, carboxy, CONR^7R^8 , $\text{SO}_2\text{NR}^7\text{R}^8$, SO_2R^4 , trifluoromethyl, NHSO_2R^4 , NHCOR^4 , ethylenedioxy, methylenedioxy, C_{1-6} alkyl, C_{1-6}
15 alkoxy, NR^7R^8 or SR^7 where R^7 and R^8 are independently hydrogen or C_{1-6} alkyl;

R and R^1 are independently a group $\text{Y}(\text{CH}_2)_p\text{R}^9$ where p is 0, 1, 2 or 3 and Y is O or NR^{10} where R^{10} is hydrogen, C_{1-6} alkyl or C_{3-6} cycloalkyl;
and R^9 is hydrogen, C_{1-6} alkyl which can optionally contain one or more O, S or NR^4
20 groups where R^4 is hydrogen or C_{1-6} alkyl, or a 3 to 7-membered saturated ring optionally containing a carbonyl group, one or more O, S or N atoms, or an aryl or heteroaryl group containing one to four heteroatoms selected from O, S or N, the saturated ring, aryl and heteroaryl groups all being optionally substituted by halogen, amino, hydroxy, cyano, nitro, carboxy, CONR^7R^8 , $\text{SO}_2\text{NR}^7\text{R}^8$, SO_2R^4 , trifluoromethyl, NHSO_2R^4 , NHCOR^4 ,
25 ethylenedioxy, methylenedioxy, C_{1-6} alkyl, C_{1-6} alkoxy, SR^5 or $\text{NR}^{11}\text{R}^{12}$ where R^{11} and R^{12} are independently hydrogen, C_{1-6} alkyl or together with the nitrogen atom to which they are attached form a 5- or 6-membered saturated ring optionally containing a further O, S or NR^4 group;

or R/R^1 is a group $\text{NR}^{10}(\text{CHR}^{10})\text{CONR}^2\text{R}^3$ or $\text{NR}^{10}(\text{CH}_2)_q\text{CONR}^2\text{R}^3$ where q is 1, 2 or 3;
30 or R/R^1 is a group $\text{NR}^{13}\text{R}^{14}$ where R^{13} and R^{14} together with the nitrogen atom to which they are attached form a 4 to 7-membered saturated ring optionally containing a carbonyl group, O, S or N atom and optionally substituted by C_{1-6} alkyl, amino, hydroxy, $\text{CO}_2\text{C}_{1-6}$ alkyl, halogen, NR^5R^6 , NR^7R^8 , C_{1-6} alkyl $\text{NR}^{17}\text{R}^{18}$ where R^{17} and R^{18} are independently hydrogen or C_{1-6} alkyl, $\text{CONR}^{15}\text{R}^{16}$ where R^{15} and R^{16} are independently hydrogen or C_{1-6}
35 alkyl, or optionally substituted by aryl, phenoxy, COPhenyl , or a heteroaryl group, the latter four groups being optionally substituted by halogen, amino, hydroxy, cyano, nitro,

carboxy, CONR^7R^8 , $\text{SO}_2\text{NR}^7\text{R}^8$, SO_2R^4 , trifluoromethyl, NHSO_2R^4 , NHCOR^4 , ethylenedioxy, methylenedioxy, C_{1-6} alkyl, C_{1-6} alkoxy, SR^5 or $\text{NR}^{11}\text{R}^{12}$ where R^{11} and R^{12} are independently hydrogen, C_{1-6} alkyl or together with the nitrogen atom to which they are attached form a 5- or 6-membered saturated ring optionally containing a further O, S or NR⁴ group;

and pharmaceutically acceptable salts or solvates thereof.

In the context of the present specification, unless otherwise indicated, an alkyl or alkenyl group or an alkyl or alkenyl moiety in a substituent group may be linear or branched. Aryl groups include phenyl and naphthyl. Heteroaryl groups include 5- or 6- membered, 5,6- or 6,6-fused heterocyclic rings containing one or more heteroatoms selected from N, S, O. Examples include pyridine, pyrimidine, thiazole, oxazole, pyrazole, imidazole, furan, thiophene, quinoline, isoquinoline, benzimidazole, benzofuran, benzothiophene and indole.

Certain compounds of formula (I) are capable of existing in stereoisomeric forms. It will be understood that the invention encompasses all geometric and optical isomers of the compounds of formula (I) and mixtures thereof including racemates. Tautomers and mixtures thereof also form an aspect of the present invention.

Preferably X is CH, NHR^2 , OR^2 where R^2 is preferably H or C_{1-6} alkyl

Preferably R is a group $\text{Y}(\text{CH}_2)_p\text{R}^7$ where p is 0 or 1 and Y is NR^8 where R^8 is hydrogen and R^7 is substituted phenyl. Preferably R^7 is phenyl substituted by halogen, especially chloro. More preferably R^7 is phenyl substituted by chloro in the 4-position.

Preferably R^1 is a group $\text{NR}^{13}\text{R}^{14}$ where R^{13} and R^{14} together with the nitrogen atom to which they are attached form a morpholine ring, piperidine or piperazine ring optionally substituted, or R^1 is a group NR^9R^{10} where R^{10} is H or C_{1-6} alkyl and R^9 is C_{1-6} alkyl which can optionally contain one or more O, S or NR⁴ groups where R^4 is hydrogen or C_{1-6} alkyl.

The most preferred substituents for R and R^1 are those of the examples exemplified herein.

Preferred compounds of the invention include:


4-[(4-Chlorophenyl)amino]-6-(dimethylamino)-1,3,5-triazine-2-carbonitrile,
4-Morpholin-4-yl-6-(4-phenoxy piperidin-1-yl)-1,3,5-triazine-2-carbonitrile,

- 4-[(4-Chlorophenyl)amino]-6-morpholin-4-yl-1,3,5-triazine-2-carbonitrile,
 4-(7-Azabicyclo[2.2.1]hept-7-yl)-6-[(4-chlorophenyl)amino]-1,3,5-triazine-2-carbonitrile,
 4-[(4-Chlorophenyl)amino]-6-pyrrolidin-1-yl-1,3,5-triazine-2-carbonitrile,
 4-[(4-Chlorophenyl)amino]-6-piperidin-1-yl-1,3,5-triazine-2-carbonitrile,
 5 4-[(4-Chlorophenyl)amino]-6-(ethylamino)-1,3,5-triazine-2-carbonitrile,
 4-[(4-Chlorophenyl)amino]-6-(3-hydroxypyrrolidin-1-yl)-1,3,5-triazine-2-carbonitrile,
 4-[(4-Chlorophenyl)amino]-6-[(2-piperidin-1-ylethyl)amino]-1,3,5-triazine-2-carbonitrile,
 4-[(4-Chlorophenyl)amino]-6-(4-phenylpiperidin-1-yl)-1,3,5-triazine-2-carbonitrile,
 4-[(3-Chlorobenzyl)amino]-6-(dimethylamino)-1,3,5-triazine-2-carbonitrile,
 10 4-Morpholin-4-yl-6-[(4-morpholin-4-ylphenyl)amino]-1,3,5-triazine-2-carbonitrile,
 4-(2,3-Dihydro-1,4-benzodioxin-6-ylamino)-6-morpholin-4-yl-1,3,5-triazine-2-carbonitrile,
 4-Morpholin-4-yl-6-(3-phenylpiperidin-1-yl)-1,3,5-triazine-2-carbonitrile,
 4-(1,4'-Bipiperidin-1'-yl)-6-morpholin-4-yl-1,3,5-triazine-2-carbonitrile,
 15 4-[4-(1H-Imidazol-1-yl)piperidin-1-yl]-6-morpholin-4-yl-1,3,5-triazine-2-carbonitrile,
 4-[4-(4-Chlorobenzoyl)piperidin-1-yl]-6-morpholin-4-yl-1,3,5-triazine-2-carbonitrile,
 4-[4-(5-Chloropyridin-2-yl)piperazin-1-yl]-6-morpholin-4-yl-1,3,5-triazine-2-carbonitrile,
 4-Morpholin-4-yl-6-[[3-(2-oxopyrrolidin-1-yl)propyl]amino]-1,3,5-triazine-2-carbonitrile,
 1-(4-Cyano-6-morpholin-4-yl-1,3,5-triazin-2-yl)-N,N-diethylpiperidine-3-carboxamide,
 20 4-[4-(2-Methoxyphenyl)piperazin-1-yl]-6-morpholin-4-yl-1,3,5-triazine-2-carbonitrile,
 N~2~(4-Cyano-6-morpholin-4-yl-1,3,5-triazin-2-yl)-N~1~,N~1~-bis{4-[N-(4-cyano-6-morpholin-4-yl-1,3,5-triazin-2-yl)-N-isobutylglycyl]morpholin-3-yl}-N~2~-isobutylglycinamide,
 4-Morpholin-4-yl-6-[(2-pyridin-3-ylethyl)amino]-1,3,5-triazine-2-carbonitrile,
 25 4-[[2-(2-Furyl)ethyl]amino]-6-morpholin-4-yl-1,3,5-triazine-2-carbonitrile,
 4-[(4-Chlorophenyl)amino]-6-(4-methylpiperazin-1-yl)-1,3,5-triazine-2-carbonitrile,
 4-Azetidin-1-yl-6-[(4-chlorophenyl)amino]-1,3,5-triazine-2-carbonitrile,
 4-[(4-Chlorophenyl)amino]-6-morpholin-4-ylpyrimidine-2-carbonitrile,
 4-[(4-Methylcyclohexyl)amino]-6-morpholin-4-ylpyrimidine-2-carbonitrile,
 30 4-(4-Chlorophenoxy)-6-morpholin-4-ylpyrimidine-2-carbonitrile,
 4-[(4-Chlorophenyl)amino]-6-(dimethylamino)pyrimidine-2-carbonitrile,
 4-[(1-Methylpiperidin-4-yl)amino]-6-morpholin-4-ylpyrimidine-2-carbonitrile,
 4-(Cyclohexylamino)-6-morpholin-4-ylpyrimidine-2-carbonitrile,
 4-[(4-Chlorophenyl)amino]-6-pyrrolidin-1-ylpyrimidine-2-carbonitrile,
 35 4-[(6-Chloropyridin-3-yl)amino]-6-morpholin-4-ylpyrimidine-2-carbonitrile,
 1-{6-[(4-Chlorophenyl)amino]-2-cyanopyrimidin-4-yl}-L-prolinamide,

- 4-(4-Aminopiperidin-1-yl)-6-[(4-chlorophenyl)amino]pyrimidine-2-carbonitrile,
 4-[(4-Chlorophenyl)amino]-6-(4-pyrrolidin-1-ylpiperidin-1-yl)pyrimidine-2-carbonitrile,
 4-[(4-Chlorophenyl)amino]-6-[(3-pyrrolidin-1-ylpropyl)amino]pyrimidine-2-carbonitrile,
 tert-Butyl 4-{6-[(4-chlorophenyl)amino]-2-cyanopyrimidin-4-yl}piperazine-1-carboxylate,
 5 4-[(4-Chlorophenyl)amino]-6-(cyclopropylamino)pyrimidine-2-carbonitrile,
 4-[(4-Chlorophenyl)amino]-6-piperazin-1-ylpyrimidine-2-carbonitrile,
 (2S)-N~2~-{6-[(4-Chlorophenyl)amino]-2-cyanopyrimidin-4-yl}-N~1~,N~1~-bis[4-(N-
 {6-[(4-chlorophenyl)amino]-2-cyanopyrimidin-4-yl}-L-leucyl)morpholin-3-yl]-L-
 leucinamide,
 10 5-Chloro-4-[(4-chlorophenyl)amino]-6-morpholin-4-ylpyrimidine-2-carbonitrile,
 4-[(4-Chlorophenyl)amino]-5-methoxy-6-piperazin-1-ylpyrimidine-2-carbonitrile,
 4-[(4-Chlorophenyl)amino]-5-methoxy-6-morpholin-4-ylpyrimidine-2-carbonitrile,
 4-[(3S)-3-Aminopyrrolidin-1-yl]-6-[(4-chlorophenyl)amino]-5-methoxypyrimidine-2-
 carbonitrile,
 15 4-[(4-Chlorophenyl)amino]-6-{4-[3-(dimethylamino)propyl]piperazin-1-yl}-5-
 methoxypyrimidine-2-carbonitrile,
 4-[(4-Chlorophenyl)amino]-6-(dimethylamino)-5-methoxypyrimidine-2-carbonitrile,
 4-[(4-Chlorophenyl)amino]-5-methoxy-6-(3-oxopiperazin-1-yl)pyrimidine-2-carbonitrile,
 1-{6-[(4-Chlorophenyl)amino]-2-cyano-5-methoxypyrimidin-4-yl}piperidine-3-
 20 carboxamide,
 4-(4-Aminopiperidin-1-yl)-6-[(4-chlorophenyl)amino]-5-methoxypyrimidine-2-
 carbonitrile,
 5-Amino-4-[(4-chlorophenyl)amino]-6-morpholin-4-ylpyrimidine-2-carbonitrile,
 5-Amino-4-[(4-Chlorophenyl)amino]-6-(ethylamino)pyrimidine-2-carbonitrile,
 25 and pharmaceutically acceptable salts thereof.

The present invention further provides a process for the preparation of a compound of formula (I) which comprises


- 30 (i) reaction of a compound of general formula (II)




L1 and L2 may be displaced by R and R¹ respectively where R and R¹ are defined in formula (I) and L3 may be displaced by a cyanide salt. The sequence of displacement of L1, L2 and L3 may be varied.

O=C1NC(=O)C(A)C(=O)N1

(III)



(IV)



(V)

(ii) when $X=CA$ and $A=NH_2$ reaction of a compound of general formula (V) under a hydrogen atmosphere with palladium catalyst at room temperature.

According to a further feature of the invention there is provided a compound of the formula (I), or a pharmaceutically acceptable salt thereof, for use as a therapeutic agent.

According to a further feature of the present invention there is provided a method for producing inhibition of a cysteine protease in a warm blooded animal, such as man, in need of such treatment, which comprises administering to said animal an effective amount of a compound of the present invention, or a pharmaceutically acceptable salt thereof.

The invention also provides a compound of the formula (I), or a pharmaceutically acceptable salt thereof, for use as a medicament; and the use of a compound of the formula

(I) of the present invention, or a pharmaceutically acceptable salt thereof, in the manufacture of a medicament for use in the inhibition of a cysteine protease in a warm blooded animal, such as man.

5 In particular the invention provides the use of a compound of the formula (I) of the present invention, or a pharmaceutically acceptable salt thereof, in the manufacture of a medicament for use in the inhibition of Cathepsin S in a warm blooded animal, such as man. In order to use a compound of the formula (I) or a pharmaceutically acceptable salt thereof for the therapeutic treatment of mammals including humans, in particular in the
10 inhibition of a cysteine protease, it is normally formulated in accordance with standard pharmaceutical practice as a pharmaceutical composition.

Therefore in another aspect the present invention provides a pharmaceutical composition which comprises a compound of the formula (I) or a pharmaceutically acceptable salt
15 thereof and a pharmaceutically acceptable diluent or carrier.

The pharmaceutical compositions of this invention may be administered in standard manner for the disease condition that it is desired to treat, for example by oral, rectal or parenteral administration. For these purposes the compounds of this invention may be
20 formulated by means known in the art into the form of, for example, tablets, capsules, aqueous or oily solutions or suspensions, (lipid) emulsions, dispersible powders, suppositories, ointments, creams, drops and sterile injectable aqueous or oily solutions or suspensions.

25 A suitable pharmaceutical composition of this invention is one suitable for oral administration in unit dosage form, for example a tablet or capsule which contains between 100 mg and 1 g of the compound of this invention.

In another aspect a pharmaceutical composition of the invention is one suitable for
30 intravenous, subcutaneous or intramuscular injection.

Each patient may receive, for example, an intravenous, subcutaneous or intramuscular dose of 1 mgkg^{-1} to 100 mgkg^{-1} of the compound, preferably in the range of 5 mgkg^{-1} to 20 mgkg^{-1} of this invention, the composition being administered 1 to 4 times per day. The
35 intravenous, subcutaneous and intramuscular dose may be given by means of a bolus injection. Alternatively the intravenous dose may be given by continuous infusion over a

period of time. Alternatively each patient will receive a daily oral dose which is approximately equivalent to the daily parenteral dose, the composition being administered 1 to 4 times per day.

- 5 The following illustrate representative pharmaceutical dosage forms containing the compound of formula (I), or a pharmaceutically-acceptable salt thereof (hereafter compound X), for therapeutic or prophylactic use in humans:

(a)

<u>Tablet I</u>	<u>mg/tablet</u>
Compound X.	100
Lactose Ph.Eur.	179
Croscarmellose sodium	12.0
Polyvinylpyrrolidone	6
Magnesium stearate	3.0

(b)

<u>Tablet II</u>	<u>mg/tablet</u>
Compound X	50
Lactose Ph.Eur.	229
Croscarmellose sodium	12.0
Polyvinylpyrrolidone	6
Magnesium stearate	3.0

s

(c)

<u>Tablet III</u>	<u>mg/tablet</u>
Compound X	1.0
Lactose Ph.Eur.	92
Croscarmellose sodium	4.0
Polyvinylpyrrolidone	2.0
Magnesium stearate	1.0

(d)

<u>Capsule</u>	<u>mg/capsule</u>
Compound X	10
Lactose Ph.Eur.	389
Croscarmellose sodium	100
Magnesium stearate	1.

(e)

<u>Injection I</u>	<u>(50 mg/ml)</u>
Compound X	5.0% w/v
Isotonic aqueous solution	to 100%

Buffers, pharmaceutically-acceptable cosolvents such as polyethylene glycol, polypropylene glycol, glycerol or ethanol or complexing agents such as hydroxy-propyl β cyclodextrin may be used to aid formulation.

5 Note

The above formulations may be obtained by conventional procedures well known in the pharmaceutical art. The tablets (a)-(c) may be enteric coated by conventional means, for example to provide a coating of cellulose acetate phthalate.

10 The following examples illustrate the invention.

Patent 3,726,000

Example 1

4-[(4-Chlorophenyl)amino]-6-(dimethylamino)-1,3,5-triazine-2-carbonitrile

(i) 4,6-Dichloro-N-(4-chlorophenyl)-1,3,5-triazin-2-amine

4-Chloroaniline (8.28g) was added to a mixture of trichlorotriazine (6g) in acetone/ice-water (1:1, 60ml) and stirred for 1h. The solid was filtered off and dried to give a light brown solid, 8.5g.

MS: APCI(+ve) 275/7(M+1)

(ii) 6-Chloro-N-2-[(4-chlorophenyl)-N-4,N-4-dimethyl-1,3,5-triazine-2,4-diamine

A solution of dimethylamine in tetrahydrofuran (2M, 1.1ml) was added to a mixture of the product from step (i) (0.3g) in acetone (10ml) and ice-water (10ml). After stirring for 1h, the solid was filtered, washed with water and dried. Yield 0.3g solid.

MS: APCI(+ve) 284(M+1)

(iii) 4-[(4-Chlorophenyl)amino]-6-(dimethylamino)-1,3,5-triazine-2-carbonitrile

Sodium cyanide (0.138g) was added to a solution of the product from step (ii) (0.4g) in N,N-dimethylformamide (20ml) and heated at 90°C for 16h. The mixture was partitioned between ethyl acetate and water, a solid formed which was filtered off and purified by RPHPLC 15-85% acetonitrile in aqueous trifluoroacetic acid. Yield 0.05g

MS: APCI(+ve) 275(M+1)

¹H NMR: (DMSO-d₆) δ 10.30(1H, bs), 7.72-7.37(4H, 2xd), 3.14(6H, s)

Example 2

4-Morpholin-4-yl-6-(4-phenoxy-piperidin-1-yl)-1,3,5-triazine-2-carbonitrile

(i) 2,4-Dichloro-6-morpholin-4-yl-1,3,5-triazine

Morpholine was added dropwise to stirred solution of trichlorotriazine (6.7g), N,N-diisopropylethylamine (6.5ml) in dichloromethane (50ml) at -78°C. The solid formed was filtered off, washed with water, dried to give a white solid (6.7g).

MS: APCI(+ve) 235(M+1)

(ii) 4-Morpholin-4-yl-6-(4-phenoxy piperidin-1-yl)-1,3,5-triazine-2-carbonitrile

4-Phenoxy piperidine (0.15g) was added to a solution of the product from step (i) (0.2g), N,N-diisopropylethylamine (1.47ml) in tetrahydrofuran and stirred at room temperature for 16h. The solvent was evaporated under reduced pressure and the residue purified by chromatography on silica eluting with ether/isohexane (1:2). Yield 0.3g white solid. The solid was dissolved in N,N-dimethylformamide (20ml), sodium cyanide (0.1g) added and heated at 90°C for 32h. The mixture was partitioned between ethyl acetate and water, the organic layer separated, washed with water, brine, dried (MgSO₄) and evaporated under reduced pressure. The residue was purified by RPHPLC 35-95% acetonitrile in aqueous trifluoroacetic acid. Yield 0.079g

MS: APCI(+ve) 367(+H)

¹H NMR: (DMSO-d₆) δ 7.31-6.91(5H, m), 4.66(1H, m), 4.11-4.04(2H, m), 3.70-3.57(10H, m), 2.00-1.95(2H, m), 1.63-1.60(2H, m)

Examples 3-26

Examples 3-26 were prepared according to the methods of example 1 or 2 using the appropriate amines.

Example 3**4-[(4-Chlorophenyl)amino]-6-morpholin-4-yl-1,3,5-triazine-2-carbonitrile**

MS: APCI(-ve) 315(M-1)

¹H NMR: (DMSO-d₆) δ 10.41(1H, bs), 7.66-7.31(4H, 2xd), 3.74-3.64(8H, m)

Example 4**4-(7-Azabicyclo[2.2.1]hept-7-yl)-6-[(4-chlorophenyl)amino]-1,3,5-triazine-2-carbonitrile**

MS: APCI(-ve) 325(M-1)

¹H NMR: (DMSO-d₆) δ 10.37(1H, bs), 7.70-7.35(4H, 2xd), 4.64-4.61(2H, m), 1.74-1.52(8H, m)

Example 5**4-[(4-Chlorophenyl)amino]-6-pyrrolidin-1-yl-1,3,5-triazine-2-carbonitrile**

5 MS: APCI(-ve) 299(M-1)

1H NMR: (DMSO-d6) δ 10.29(1H, bs), 7.75-7.36(4H, 2xd), 3.54-3.49(4H, m), 1.96-1.90(4H, m)

Example 6

10 **4-[(4-Chlorophenyl)amino]-6-piperidin-1-yl-1,3,5-triazine-2-carbonitrile**

MS: APCI(-ve) 313(M-1)

1H NMR: (DMSO-d6) δ 10.29(1H, bs), 7.66-7.38(4H, 2xd), 3.74-3.73(4H, m), 1.64-1.55(6H, m)

15

Example 7**4-[(4-Chlorophenyl)amino]-6-(ethylamino)-1,3,5-triazine-2-carbonitrile**

MS: APCI(-ve) 273(M-1)

20 1H NMR: (DMSO-d6) δ 10.31(1H, bs), 8.39-7.34(5H, m), 3.36-3.27(2H, q), 1.10(3H, t)

Example 8**4-[(4-Chlorophenyl)amino]-6-(3-hydroxypyrrolidin-1-yl)-1,3,5-triazine-2-carbonitrile**

25 MS: APCI(-ve) 315(M-1)

1H NMR: (DMSO-d6) δ 10.31(1H, s), 7.76-7.38 (4H, m), 5.06(1H, m), 4.38(1H, m), 3.66-3.45(4H, m), 2.04-1.95(2H, m)

Example 9

30 **4-[(4-Chlorophenyl)amino]-6-[(2-piperidin-1-ylethyl)amino]-1,3,5-triazine-2-carbonitrile**

MS: APCI(+ve) 358(M+1)

¹H NMR: (DMSO-d₆) δ 10.38(1H, s), 8.35-7.34(5H, m), 3.37(2H, m), 2.43-2.34(6H, m), 1.48-1.44(6H, m)

5

Example 10

4-[(4-Chlorophenyl)amino]-6-(4-phenylpiperidin-1-yl)-1,3,5-triazine-2-carbonitrile

MS: APCI(-ve) 389(M-1)

10 ¹H NMR: (DMSO-d₆) δ 10.33(1H, bs), 7.68-7.16(9H, m), 4.72-4.69(2H, d), 3.09-2.85(3H, m), 1.88-1.55(4H, m)

Example 11

4-[(3-Chlorobenzyl)amino]-6-(dimethylamino)-1,3,5-triazine-2-carbonitrile

15

MS: APCI(-ve) 287(M-1)

¹H NMR: (DMSO-d₆) δ 8.59-8.44(1H, t), 7.37-7.22(4H, m), 4.47-4.44(2H, m), 3.07-3.03(6H, m)

20

Example 12

4-Morpholin-4-yl-6-[(4-morpholin-4-ylphenyl)amino]-1,3,5-triazine-2-carbonitrile

MS: APCI(+ve) 368(M+1)

25 ¹H NMR: (DMSO-d₆) δ 10.09(1H, s), 7.57-7.48(2H, d), 6.93-6.84(2H, d), 3.72-3.55(12H, m), 3.07-3.00(4H, m)

Example 13

4-(2,3-Dihydro-1,4-benzodioxin-6-ylamino)-6-morpholin-4-yl-1,3,5-triazine-2-carbonitrile

30

MS: APCI(-ve) 339(M-1)

¹H NMR: (DMSO-d₆) δ 10.13(1H, s), 7.21-6.80(3H, m), 4.23-3.64(12H, m)

Example 14

5 4-Morpholin-4-yl-6-(3-phenylpiperidin-1-yl)-1,3,5-triazine-2-carbonitrile

MS: APCI(+ve) 351(M+1)

¹H NMR: (DMSO-d₆) δ 7.35-7.22(5H, m), 4.67-2.64(13H, m), 1.93-1.48(4H, m)

10 Example 15

4-(1,4'-Bipiperidin-1'-yl)-6-morpholin-4-yl-1,3,5-triazine-2-carbonitrile, trifluoroacetate salt

MS: APCI(+ve) 358(M+1)

15 ¹H NMR: (DMSO-d₆) δ 9.23(1H, bm), 4.75-4.64(2H, m), 3.71-3.36(11H, m), 2.93-
2.87(4H, m), 2.08-1.37(10H, m)

Example 16

4-[4-(1H-Imidazol-1-yl)piperidin-1-yl]-6-morpholin-4-yl-1,3,5-triazine-2-carbonitrile

20

MS: APCI(+ve) 341(M+1)

¹H NMR: (DMSO-d₆) δ 9.11(1H, s), 7.89-7.88(1H, s), 7.69-7.68(1H, s), 4.79-4.62(3H, m), 3.74-3.02(10H, m), 2.17-1.88(4H, m)

25 Example 17

4-[4-(4-Chlorobenzoyl)piperidin-1-yl]-6-morpholin-4-yl-1,3,5-triazine-2-carbonitrile

MS: APCI(+ve) 413(M+1)

30 ¹H NMR: (DMSO-d₆) δ 8.04-8.02(2H, d), 7.64-7.60(2H, d), 4.62-4.52(2H, m), 3.80-
3.69(5H, m), 3.32-3.08(6H, m), 1.87-1.43(4H, m)

Example 18**4-[4-(5-Chloropyridin-2-yl)piperazin-1-yl]-6-morpholin-4-yl-1,3,5-triazine-2-carbonitrile**

5 MS: APCI(+ve) 387(M+1)
1H NMR: (DMSO-d6) δ 8.14-8.13(1H, s), 7.65-7.62(1H, d), 6.91-6.89(1H, d), 3.82-3.57(16H, m)

Example 19

10 **4-Morpholin-4-yl-6-[[3-(2-oxopyrrolidin-1-yl)propyl]amino]-1,3,5-triazine-2-carbonitrile**

MS: APCI(+ve) 332(M+1)
1H NMR: (DMSO-d6) δ 8.10-8.07(1H, t), 3.70-3.62(8H, m), 3.39-3.17(6H, m), 2.22-
15 1.62(6H, m)

Example 20**1-(4-Cyano-6-morpholin-4-yl-1,3,5-triazin-2-yl)-N,N-diethylpiperidine-3-carboxamide**

20 MS: APCI(+ve) 374(M+1)
1H NMR: (DMSO-d6) δ 4.39(2H, m), 3.78-3.60(4H, m), 3.33-2.63(11H, m), 1.80-1.43(4H, m), 1.16-0.92(6H, m)

Example 21

25 **4-[4-(2-Methoxyphenyl)piperazin-1-yl]-6-morpholin-4-yl-1,3,5-triazine-2-carbonitrile**

MS: APCI(+ve) 382(M+1)
1H NMR: (DMSO-d6) δ 6.99-6.87(4H, m), 3.86-3.63(15H, m), 2.99(4H, m)

Example 22

N~2~(4-Cyano-6-morpholin-4-yl-1,3,5-triazin-2-yl)-N~1~,N~1~-bis{4-[N-(4-cyano-6-morpholin-4-yl-1,3,5-triazin-2-yl)-N-isobutylglycyl]morpholin-3-yl}-N~2~isobutylglycinamide

MS: APCI(+ve) 390(M+1)

¹H NMR: (DMSO-d₆) δ 4.39-4.36(2H, d), 3.62-3.31(18H, m), 2.05-1.92(1H, m), 0.87-0.85(6H, d)

Example 23

4-Morpholin-4-yl-6-[(2-pyridin-3-ylethyl)amino]-1,3,5-triazine-2-carbonitrile, trifluoroacetate salt

MS: APCI(+ve) 312(M+1)

¹H NMR: (DMSO-d₆) δ 8.69-8.64(2H, m), 8.22-8.04(2H, m), 7.78-7.70(1H, m), 3.66-3.53(10H, m), 2.97-2.93(2H, t)

Example 24

4-[(2-(2-Furyl)ethyl)amino]-6-morpholin-4-yl-1,3,5-triazine-2-carbonitrile

MS: APCI(-ve) 299(M+1)

¹H NMR: (DMSO-d₆) δ 8.21(1H, t), 7.51(1H, s), 6.34(1H, s), 6.15(1H, s), 3.64-3.62(8H, m), 3.52-3.46(2H, m), 2.86-2.82(2H, m)

Example 25

4-[(4-Chlorophenyl)amino]-6-(4-methylpiperazin-1-yl)-1,3,5-triazine-2-carbonitrile, trifluoroacetate salt

MS: APCI(+ve) 330(M+1)

¹H NMR: (DMSO-d₆) δ 10.56(1H, bs), 10.05(1H, brs), 7.66-7.40(4H, m), 3.41-3.35(8H, m), 2.81(3H, s)

Example 26**4-Azetidin-1-yl-6-[(4-chlorophenyl)amino]-1,3,5-triazine-2-carbonitrile**

MS: APCI(-ve) 285(M-1)

1H NMR: (DMSO-d₆) δ 10.32(1H, s), 7.72-7.32(4H, m), 4.15-4.10(4H, m), 2.38-2.30(2H, q)

Example 27**4-[(4-Chlorophenyl)amino]-6-morpholin-4-ylpyrimidine-2-carbonitrile****(i) N-(4-Chlorophenyl)-2,6-difluoropyrimidin-4-amine**

4-Chloroaniline was added to a stirred solution of 2,4,6- trifluoropyrimidine (7.7g), potassium carbonate (7.86g) in ethanol (80ml). The mixture was stirred at room temperature for 16h, diluted with water, extracted with ethyl acetate, dried (MgSO₄) and evaporated under reduced pressure. The residue was purified by chromatography on silica eluting with isohexane/ethyl acetate (4:1). Yield 8.3g cream solid

1H NMR: (DMSO-d₆) δ 10.47(1H, s), 7.58(2H, d), 7.45(2H, d), 6.35(1H, s)

(ii) 4-[(4-Chlorophenyl)amino]-6-fluoropyrimidine-2-carbonitrile

Sodium cyanide (0.046g) was added to a solution of the product from step (i) (0.113g) in dimethylsulphoxide (3ml) and stirred at room temperature for 1.5h. The mixture was partitioned between ethyl acetate and water, the organics washed with water, dried (MgSO₄), and evaporated under reduced pressure. The residue was purified by chromatography on silica eluting with isohexane/ethyl acetate (4:1). Yield 0.036g

1H NMR: (DMSO-d₆) δ 10.56(1H, s), 7.57(2H, d), 7.47(2H, d), 6.65(1H, s)

(iii) 4-[(4-Chlorophenyl)amino]-6-morpholin-4-ylpyrimidine-2-carbonitrile

Morpholine (0.16g) was added to a solution of the product from step (ii) (0.16g) in isopropylalcohol (4ml) and stirred for 2h at room temperature. The mixture was partitioned between ethyl acetate and aqueous sodium hydrogencarbonate solution, the organics separated, dried (MgSO₄) and evaporated under reduced pressure. The residue was purified by chromatography on silica eluting with isohexane/ethyl acetate (1:1). Yield 0.09g

MS: APCI(+ve) 316(M+1)

¹H NMR: (DMSO-d₆) δ 9.65(1H, s), 7.52(2H, d), 7.38(2H, d), 6.08(1H, s), 3.67(4H, t), 3.48(2H, t)

5 **Examples 28-42**

Examples 28-42 were prepared according to the general method of example 27 using the appropriate amines or phenols

Example 28

10 **4-[(4-Methylcyclohexyl)amino]-6-morpholin-4-ylpyrimidine-2-carbonitrile**

MS: APCI(+ve) 302(M+1)

¹H NMR: (DMSO-d₆) δ 7.21-7.18(1H, d), 5.83(1H, s), 3.89(1H, bs), 3.88(4H, m), 3.41(4H, m), 1.63-1.28(9H, m), 0.89(3H, d)

15

Example 29

4-(4-Chlorophenoxy)-6-morpholin-4-ylpyrimidine-2-carbonitrile

MS: APCI(-ve) 315(M-1)

20 ¹H NMR: (DMSO-d₆) δ 7.53-7.20(4H, 2xd), 6.63(1H, s), 3.65-3.63(8H, m)

Example 30

4-[(4-Chlorophenyl)amino]-6-(dimethylamino)pyrimidine-2-carbonitrile

25 MS: APCI(+ve) 274(M+1)

¹H NMR: (DMSO-d₆) δ 9.57(1H, s), 7.55-7.34(4H, 2xd), 5.93(1H, s), 3.02(6H, m)

Example 31

30 **4-[(1-Methylpiperidin-4-yl)amino]-6-morpholin-4-ylpyrimidine-2-carbonitrile, trifluoroacetate salt**

MS: APCI(+ve) 303(M+1)

¹H NMR: (DMSO-d₆) δ 9.36(1H, brs), 7.49-7.47(1H, d), 5.76(1H, s), 3.92(1H, bm), 3.67-3.43(8H, 2xm), 3.34-3.10(4H, m), 2.75(3H, s), 2.07-1.68(4H, m)

5 **Example 32**

4-(Cyclohexylamino)-6-morpholin-4-ylpyrimidine-2-carbonitrile

MS: APCI(+ve) 288(M+1)

¹H NMR: (DMSO-d₆) δ 7.23-7.21(1H, d), 5.73(1H, s), 3.62-3.42(9H, m), 1.83-1.07(10H, m)

Example 33

4-[(4-Chlorophenyl)amino]-6-pyrrolidin-1-ylpyrimidine-2-carbonitrile

15 MS: APCI(+ve) 300(M+1)

¹H NMR: (DMSO-d₆) δ 9.55(1H, s), 7.54-7.35(4H, 2xd), 5.79(1H, s), 3.38(4H, m), 1.93(4H, m)

Example 34

20 **4-[(6-Chloropyridin-3-yl)amino]-6-morpholin-4-ylpyrimidine-2-carbonitrile**

MS: APCI(-ve) 315(M-1)

¹H NMR: (DMSO-d₆) δ 9.83(1H, s), 8.55-8.49(1H, s), 8.06-8.02(1H, d), 7.49-7.46(1H, d), 6.10(1H, s), 3.69-3.66(4H, m), 3.52-3.48(4H, m)

25 **Example 35**

1-{6-[(4-Chlorophenyl)amino]-2-cyanopyrimidin-4-yl}-L-prolinamide

MS: APCI(+ve) 343(M+1)

30 ¹H NMR: (DMSO-d₆) δ 9.33(1H, s), 7.57-7.24(4H, 2xd), 7.00(2H, bm), 5.81(1H, s), 4.31-3.38(3H, m), 2.26-1.26(4H, m)

Example 36

4-(4-Aminopiperidin-1-yl)-6-[(4-chlorophenyl)amino]pyrimidine-2-carbonitrile, trifluoroacetate salt

MS: APCI(+ve) 329(M+1)

¹H NMR: (DMSO-d₆) δ 9.66(1H, s), 7.88-7.36(7H, m), 6.15(1H, s), 4.23-4.20(2H, m), 3.17-2.96(3H, m), 1.98-1.38(4H, m)

Example 37

4-[(4-Chlorophenyl)amino]-6-(4-pyrrolidin-1-ylpiperidin-1-yl)pyrimidine-2-carbonitrile, acetate salt

MS: APCI(+ve) 383(M+1)

¹H NMR: (DMSO-d₆) δ 9.57(1H, s), 7.53-7.35(4H, 2xd), 6.09(1H, s), 4.06-2.51(9H, m), 1.92-1.90(5H, m), 1.74-1.34(6H, m)

Example 38

4-[(4-Chlorophenyl)amino]-6-[(3-pyrrolidin-1-ylpropyl)amino]pyrimidine-2-carbonitrile, trifluoroacetate salt

MS: APCI(+ve) 357(M+1)

¹H NMR: (DMSO-d₆) δ 9.52(2H, m), 7.63-7.36(5H, 2xd+m), 5.92(1H, bs), 3.54-2.99(8H, m), 2.00-1.84(6H, m)

Example 39

tert-Butyl 4-{6-[(4-chlorophenyl)amino]-2-cyanopyrimidin-4-yl}piperazine-1-carboxylate

MS: APCI(+ve) 415(M+1)

¹H NMR: (DMSO-d₆) δ 9.66(1H, s), 7.52(2H, d), 7.37(2H, d), 6.07(1H, s), 3.54-3.51(4H, m), 3.44-3.41(4H, m), 1.42(9H, s)

Example 40

4-[(4-Chlorophenyl)amino]-6-(cyclopropylamino)pyrimidine-2-carbonitrile

MS: APCI(+ve) 286(M+1)

5 1H NMR: (DMSO-d₆) δ 9.65(1H, s), 7.80(1H, s), 7.53(2H, d), 7.37(2H, d), 6.08(1H, s),
0.76-0.71(2H, m), 0.50-0.46(2H, m)

Example 41

4-[(4-Chlorophenyl)amino]-6-piperazin-1-ylpyrimidine-2-carbonitrile

10

MS: APCI(+ve) 315(M+1)

1H NMR: (DMSO-d₆) δ 9.62(1H, s), 7.53(2H, d), 7.37(2H, d), 6.07(1H, s), 3.46(4H, t),
2.79(4H, t)

15

Example 42

**(2S)-N~2~-{6-[(4-Chlorophenyl)amino]-2-cyanopyrimidin-4-yl}-N~1~,N~1~-bis[4-(N-
{6-[(4-chlorophenyl)amino]-2-cyanopyrimidin-4-yl}-L-leucyl)morpholin-3-yl]-L-
leucinamide**

20

MS: APCI(+ve) 429(M+1)

1H NMR: (DMSO-d₆) δ 9.51(1H, s), 7.82(1H, d), 7.46(2H, d), 7.37(2H, d), 6.09(1H, s),
4.87(1H, s), 3.67-3.47(6H, m), 3.35-3.25(2H, m), 1.66-1.53(2H, m), 1.48-1.39(1H, m),
0.92-0.89(6H, m)

25

Example 43

5-Chloro-4-[(4-chlorophenyl)amino]-6-morpholin-4-ylpyrimidine-2-carbonitrile

(i) 4-(5-Chloro-2,6-difluoropyrimidin-4-yl)morpholine

30

Morpholine (0.774mg) was added to a solution of 5-chloro-2,4,6-trifluoropyrimidine
(1.5g), N,N-diisopropylethylamine (1.15g) in 1,4-dioxane (30ml) and stirred at room
temperature for 16h. The mixture was partitioned between ethyl acetate and water, the
organic layer dried (MgSO₄) and evaporated under reduced pressure. The residue was

purified by chromatography on silica eluting with 8% ethyl acetate in isohexane. Yield 0.88g

(ii) 5-Chloro-N-(4-chlorophenyl)-2-fluoro-6-morpholin-4-ylpyrimidin-4-amine

4-Chloroaniline (1.44g) was added to a solution of the product from step (i) (0.88g) and N,N-diisopropylethylamine (0.484g) in 1,4-dioxane (15ml) and isopropylalcohol (15ml) and the mixture heated at 110°C for 6 days. The mixture was partitioned between ethyl acetate and water, the organics dried (MgSO₄), and evaporated under reduced pressure. The solid was triturated with ethyl acetate, filtered and the filtrate purified by chromatography on silica eluting with 3% ethyl acetate in toluene. Yield 0.28g

MS: APCI(+ve) 343/5(M+1)

(iii) 5-Chloro-4-[(4-chlorophenyl)amino]-6-morpholin-4-ylpyrimidine-2-carbonitrile

Sodium cyanide (0.057g) was added to a solution of the product from step (ii) (0.2g) in dimethylsulphoxide (5ml) and the mixture stirred at room temperature. After 18h, the mixture was partitioned between ethyl acetate and water, the organics separated, washed with water, dried (MgSO₄) and evaporated under reduced pressure. The residue was purified by chromatography on silica eluting with 5% ethyl acetate in toluene. Yield 0.09g

MS: APCI(-ve) 348(M-1)

¹H NMR: (DMSO-d₆) δ 9.34(1H, s), 7.53(2H, d), 7.42(2H, d), 3.71(4H, t), 3.55(4H, t)

Example 44

4-[(4-Chlorophenyl)amino]-5-methoxy-6-piperazin-1-ylpyrimidine-2-carbonitrile, trifluoroacetate salt

(i) 5-Methoxy-2-thioxodihydropyrimidine-4,6(1H,5H)-dione

Thiourea (24g) and methoxymethyl malonate (34g) was added to a solution of sodium (12g) in methanol and the mixture heated under reflux for 10h. The methanol was evaporated under reduced pressure, water (500ml) added and extracted with ether. The aqueous layer was acidified to pH1 with conc. hydrochloric acid, evaporated to ~200ml and the precipitate filtered and dried. Yield 23g

¹H NMR: (DMSO-d₆) δ 11.31(2H, s), 3.48(3H, s)

(ii) 2-(Ethylthio)-5-methoxypyrimidine-4,6(1H,5H)-dione

Ethyl iodide (11.2ml) was added dropwise to a stirred mixture of the product from step (i) (23g) and sodium hydroxide (6g) in water (400ml). After 16h the mixture was filtered, the filtrate acidified to pH1 and the precipitate filtered, washed with water and dried. Yield 17.8g

¹H NMR: (DMSO-d₆) δ 12.25(1H, s), 3.59(3H, s), 3.57(1H, s), 3.06(2H, q), 1.28(3H, t)

(iii) 4,6-Dichloro-2-(ethylthio)-5-methoxypyrimidine

A mixture of the product from step (ii) (17.8g) and N,N-diethylaniline (20ml) in phosphorus oxychloride (400ml) was heated at 100°C for 3h. The excess reagent was removed under reduced pressure and the residue poured onto ice and extracted with ether. The ether layer was washed with water, dried (MgSO₄) and evaporated under reduced pressure. The residue was purified by chromatography on silica eluting with 5% ethyl acetate in isohexane. Yield 12.8g

(iv) 6-Chloro-N-(4-chlorophenyl)-2-(ethylthio)-5-methoxypyrimidin-4-amine

A solution of the product from step (iii) (4g) and 4-chloroaniline (5.3g) in ethanol (40ml) was heated under reflux for 16h then the solvent removed under reduced pressure. The residue was partitioned between ethyl acetate and 2M hydrochloric acid, the organics washed with water, dried (MgSO₄) and evaporated under reduced pressure. The residue was purified by chromatography on silica eluting with 10-15% ethyl acetate in isohexane. Yield 4.99 g

MS: APCI(+ve) 330/2(M+1)

(v) 4-Chloro-6-[(4-chlorophenyl)amino]-5-methoxypyrimidine-2-carbonitrile

A mixture of the product from step (iv) (4.9g) and 3-chloroperoxybenzoic acid (10g, Aldrich 77% max.) in dichloromethane (150ml) was stirred at room temperature for 3h, washed with aqueous sodium metabisulphite solution, water, aqueous sodium hydrogencarbonate solution, water, dried (MgSO₄) and evaporated under reduced pressure. The solid was dissolved in dimethylsulphoxide (40ml), sodium cyanide (1.1g) added and stirred for 2h at room temperature. The mixture was partitioned between ethyl acetate and water, the organics dried (MgSO₄) and evaporated under reduced pressure. The residue

was purified by chromatography on silica eluting with 30% ethyl acetate in isohexane.

Yield 3.23g

MS: APCI(+ve) 295/7(M+1)

(vi) 4-[(4-Chlorophenyl)amino]-5-methoxy-6-piperazin-1-ylpyrimidine-2-carbonitrile

A solution of the product from step (v) (0.25g) and piperazine (0.366g) in tetrahydrofuran (8ml) was heated at 60°C for 6h then the solvent removed under reduced pressure. The residue was purified by RPHPLC 15-75% acetonitrile in aqueous trifluoroacetic acid.

Yield 0.139g

MS: APCI(+ve) 345(M+1)

¹H NMR: (DMSO-d₆) δ 9.29(1H, s), 8.92(2H, s), 7.67(2H, d), 7.40(2H, d), 3.83-3.80(4H, m), 3.69(3H, s), 3.25-3.23(4H, m)

Mpt 230°C

Examples 45-51

Examples 45-51 were prepared according to the method of example 44 using the appropriate amines

Example 45

4-[(4-Chlorophenyl)amino]-5-methoxy-6-morpholin-4-ylpyrimidine-2-carbonitrile

MS: APCI(+ve) 346(M+1)

¹H NMR: (DMSO-d₆) δ 9.20(1H, s), 7.67(2H, d), 7.38(2H, d), 3.73-3.70(4H, m), 3.68(3H, s), 3.63-3.61(4H, m)

Mpt 176°C

Example 46

4-[(3S)-3-Aminopyrrolidin-1-yl]-6-[(4-chlorophenyl)amino]-5-methoxypyrimidine-2-carbonitrile, trifluoroacetate salt

MS: APCI(+ve) 345/7(M+1)

¹H NMR: (DMSO-d₆) δ 9.12(1H, s), 8.10(3H, s), 7.66(2H, d), 7.38(2H, d), 3.91-3.69(5H, m), 3.65(3H, s), 2.33-2.22(1H, m), 2.08-2.01(1H, m)
Mpt 345-7°C

5 Example 47

4-[(4-Chlorophenyl)amino]-6-{4-[3-(dimethylamino)propyl]piperazin-1-yl}-5-methoxypyrimidine-2-carbonitrile, bis-trifluoroacetate salt

MS: APCI(+ve) 430/2(M+1)

10 ¹H NMR: (DMSO-d₆) 90°C δ 8.94(1H, s), 7.64(2H, d), 7.35(2H, d), 3.81(4H, brs),
3.70(3H, s), 3.15-3.09(2H, m), 3.00(4H, brs), 2.86(2H, brs), 2.81(6H, s), 2.03-1.95(2H, m)
Mpt 210-2°C

Example 48

15 **4-[(4-Chlorophenyl)amino]-6-(dimethylamino)-5-methoxypyrimidine-2-carbonitrile**

MS: APCI(-ve) 302/4(M-1)

1H NMR: (DMSO-d6) 90°C δ 9.08(1H, s), 7.66(2H, d), 7.37(2H, d), 3.62(3H, s), 3.13(6H, s)
Mpt 173°C

Example 49

4-[(4-Chlorophenyl)amino]-5-methoxy-6-(3-oxopiperazin-1-yl)pyrimidine-2-carbonitrile

MS: APCI(-ve) 357/9(M-1)

¹H NMR: (DMSO-d₆) δ 9.24(1H, s), 8.10(1H, s), 7.67(2H, d), 7.39(2H, d), 4.17(2H, s), 3.85-3.83(2H, m), 3.66(3H, s), 3.32-3.29(2H, m)
Mpt 244°C

Example 50**1-{6-[(4-Chlorophenyl)amino]-2-cyano-5-methoxypyrimidin-4-yl}piperidine-3-carboxamide**

- 5 MS: APCI(+ve) 387/9 (M+1)
1H NMR: (DMSO-d6) δ 9.16(1H, s), 7.68(2H, d), 7.40-7.35(3H, m), 6.89(1H, s), 4.34-4.25(2H, m), 3.65(3H, s), 3.07-2.92(2H, m), 2.35-2.40(1H, m), 1.90-1.51(4H, m)

Example 51

- 10 **4-(4-Aminopiperidin-1-yl)-6-[(4-chlorophenyl)amino]-5-methoxypyrimidine-2-carbonitrile, trifluoroacetate salt**

- MS: APCI(+ve) 359/61 (M+1)
1H NMR: (DMSO-d6) δ 9.20(1H, s), 7.93(3H, s), 7.67(2H, d), 7.39(2H, d), 4.36-4.32(2H, m), 3.66(3H, s), 3.08-3.01(2H, m), 2.00-1.97(2H, m), 1.57-1.54(2H, m)
- 15

Example 52**5-Amino-4-[(4-chlorophenyl)amino]-6-morpholin-4-ylpyrimidine-2-carbonitrile**

- 20 **(i) N-(4-Chlorophenyl)-6-morpholin-4-yl-5-nitro-2-(propylthio)pyrimidin-4-amine**
Morpholine (1.31ml) was added dropwise to a stirred solution of 4,6-dichloro-5-nitro-2-thiopropyl pyrimidine (4g), N,N-diisopropylamine (7ml) in dichloromethane (50ml) at 0°C. After 1h, 4-chloroaniline (1.9g) was added, the mixture stirred at room temperature for 24h, then heated under reflux for 24h. The mixture was partitioned between
25 dichloromethane and 2M hydrochloric acid, the organics washed with water, dried (MgSO4) and evaporated under reduced pressure. Yield 5g

MS: APCI(+ve) 410/2 (M+1)

- 30 **(ii) 4-[(4-Chlorophenyl)amino]-6-morpholin-4-yl-5-nitropyrimidine-2-carbonitrile**
A mixture of the product from step (i) (5g) and 3-chloroperoxybenzoic acid (12g, Aldrich 77% max.) in dichloromethane (200ml) was stirred at room temperature for 2h, washed with aqueous sodium metabisulphite solution, water, aqueous sodium hydrogencarbonate solution, water, dried (MgSO4) and evaporated under reduced pressure. The solid was

dissolved in dimethylsulphoxide (30ml), sodium cyanide (2g) added and stirred for 1h at room temperature. Water (500ml) was added and the solid filtered, washed with water, dried and the residue triturated with ether. Yield 1.7g

5 MS: APCI(+ve) 361/3 (M+1)

(iii) 5-Amino-4-[(4-chlorophenyl)amino]-6-morpholin-4-ylpyrimidine-2-carbonitrile

The product from step (ii) (1.7g) and 10% palladium on charcoal (0.2g) in ethyl acetate (300ml) was hydrogenated at 2Bar for 8h, filtered through celite and the solvent
10 evaporated under reduced pressure. Yield 1.05g.

MS: APCI(+ve) 329/331 (M+1)

¹H NMR: (DMSO-d₆) δ 8.66(1H, s), 7.62(2H, d), 7.39(2H, d), 5.53(2H, s), 3.78-3.76(4H, m), 3.08-3.06(4H, m)

15 Mpt 253-4°C

Example 53


5-Amino-4-[(4-Chlorophenyl)amino]-6-(ethylamino)pyrimidine-2-carbonitrile

Example 53 was prepared according to the general method of example 52 using the
20 appropriate amine

MS: APCI(+ve) 289/91(M+1)

¹H NMR: (DMSO-d₆) δ 8.19(1H, s), 7.50(2H, d), 7.31(2H, d), 6.52(1H, t), 5.20(2H, s), 3.41-3.35(2H, m), 1.18(3H, t)

25 Mpt 211-2°C



Measurement of Cathepsin S activity.


QFRET Technology (Quenched Fluorescent Resonance Energy Transfer) was used to measure the inhibition by test compounds of Cathepsin S-mediated cleavage of the synthetic peptide Z-Val-Val-Arg-AMC. Compounds were screened at five concentrations in duplicate and the pIC₅₀ values reported.

Synthetic substrate, 20μM [final]Z-Val-Val-Arg-AMC in phosphate buffer were added to a 96 well black Optiplat. The assay plates were pre-read for compound auto fluorescence on SpectraMax Gemini at 355nM excitation and 460nM emission. 250pM [final] rHuman Cathepsin S in phosphate buffer was added and incubated for 2h at room temperature on the SpectraMax Gemini, taking readings every 20min at 355nM excitation and 460nM emission.

Activity Based template (SPTB-8) used the auto fluorescent corrected data to calculate the percentage inhibition for each compound concentration using the relevant plate controls. This data was used to construct inhibition curves and pIC₅₀ estimated by non-linear regression using a 4 parameter logistic model.

CLAIMS

1. A compound of formula (I) or a salt or solvate thereof:



(I)

in which:

- 10 X is N or CA where A is hydrogen, halogen, CHR^2R^3 , OR^2 , NR^2R^3 , SR^2 ;

- 15 R^2 and R^3 are independently hydrogen, C_{1-6} alkyl or C_{3-6} cycloalkyl both of which can optionally contain one or more O, S or NR^4 groups where R^4 is hydrogen or C_{1-6} alkyl, and can be optionally substituted by aryl, heteroaryl, NR^5R^6 where R^5 and R^6 together with the nitrogen atom to which they are attached form a 4-7 membered ring optionally containing a further O, S, NR^4 , or R2 and R3 together with the nitrogen atom to which they are attached form a 4-7 membered ring optionally containing a further O, S, NR^4 group, or R2 and R3 are aryl or heteroaryl groups.

- 20 Aryl or heteroaryl groups are both optionally substituted by halogen, amino, hydroxy, cyano, nitro, carboxy, CONR^7R^8 , $\text{SO}_2\text{NR}^7\text{R}^8$, SO_2R^4 , trifluoromethyl, NHSO_2R^4 , NHCOR^4 , ethylenedioxy, methylenedioxy, C_{1-6} alkyl, C_{1-6} alkoxy, NR^7R^8 or SR^7 where R^7 and R^8 are independently hydrogen or C_{1-6} alkyl;

- 25 R and R^1 are independently a group $\text{Y}(\text{CH}_2)_p\text{R}^9$ where p is 0, 1, 2 or 3 and Y is O or NR^{10} where R^{10} is hydrogen, C_{1-6} alkyl or C_{3-6} cycloalkyl; and R^9 is hydrogen, C_{1-6} alkyl which can optionally contain one or more O, S or NR^4 groups where R^4 is hydrogen or C_{1-6} alkyl, or a 3 to 7-membered saturated ring optionally containing a carbonyl group, one or more O, S or N atoms, or an aryl or heteroaryl group containing one to four heteroatoms selected from O, S or N, the saturated ring, aryl and heteroaryl groups all being optionally substituted by halogen, amino, hydroxy, cyano, nitro, carboxy, CONR^7R^8 , $\text{SO}_2\text{NR}^7\text{R}^8$, SO_2R^4 , trifluoromethyl, NHSO_2R^4 , NHCOR^4 ,
- 30

ethylenedioxy, methylenedioxy, C₁₋₆ alkyl, C₁₋₆ alkoxy, SR⁵ or NR¹¹R¹² where R¹¹ and R¹² are independently hydrogen, C₁₋₆ alkyl or together with the nitrogen atom to which they are attached form a 5- or 6-membered saturated ring optionally containing a further O, S or NR⁴ group;

- 5 or R/R¹ is a group NR¹⁰(CHR¹⁰) CONR²R³ or NR¹⁰(CH₂)_qCONR²R³ where q is 1, 2 or 3;
 or R/R¹ is a group NR¹³R¹⁴ where R¹³ and R¹⁴ together with the nitrogen atom to which they are attached form a 4 to 7-membered saturated ring optionally containing a carbonyl group, O, S or N atom and optionally substituted by C₁₋₆ alkyl, amino, hydroxy, CO₂C₁₋₆ alkyl, halogen, NR⁵R⁶, NR⁷R⁸, C₁₋₆ alkylNR¹⁷R¹⁸ where R¹⁷ and R¹⁸ are independently
 10 hydrogen or C₁₋₆ alkyl, CONR¹⁵R¹⁶ where R¹⁵ and R¹⁶ are independently hydrogen or C₁₋₆ alkyl, or optionally substituted by aryl, phenoxy, COphenyl, or a heteroaryl group, the latter four groups being optionally substituted by halogen, amino, hydroxy, cyano, nitro, carboxy, CONR⁷R⁸, SO₂NR⁷R⁸, SO₂R⁴, trifluoromethyl, NHCO₂R⁴, NHCOR⁴, ethylenedioxy, methylenedioxy, C₁₋₆ alkyl, C₁₋₆ alkoxy, SR⁵ or NR¹¹R¹² where R¹¹ and R¹²
 15 are independently hydrogen, C₁₋₆ alkyl or together with the nitrogen atom to which they are attached form a 5- or 6-membered saturated ring optionally containing a further O, S or NR⁴ group;

and pharmaceutically acceptable salts or solvates thereof.

20.

2. A compound according to claim 1 in which X is CH, NHR², OR² where R² is hydrogen or C₁₋₆ alkyl.

25

3. A compound according to claim 1 or 2 in which R is a group Y(CH₂)_pR⁷ where p is 0 or 1 and Y is NR⁸ where R⁸ is hydrogen and R⁷ is substituted phenyl.

30

4. A compound according to any one of claims 1 to 3 in which R¹ is a group NR¹³R¹⁴ where R¹³ and R¹⁴ together with the nitrogen atom to which they are attached form a morpholine ring, piperidine or piperazine ring optionally substituted.

35

5. A compound according to any one of claims 1 to 3 in which R¹ is a group NR⁹R¹⁰ where R¹⁰ is H or C₁₋₆ alkyl and R⁹ is C₁₋₆ alkyl which can optionally contain one or more O, S or NR⁴ groups where R⁴ is hydrogen or C₁₋₆ alkyl.

6. A compound of formula (I) selected from:

4-[(4-Chlorophenyl)amino]-6-(dimethylamino)-1,3,5-triazine-2-carbonitrile,

- 4-Morpholin-4-yl-6-(4-phenoxy piperidin-1-yl)-1,3,5-triazine-2-carbonitrile,
 4-[(4-Chlorophenyl)amino]-6-morpholin-4-yl-1,3,5-triazine-2-carbonitrile,
 4-(7-Azabicyclo[2.2.1]hept-7-yl)-6-[(4-chlorophenyl)amino]-1,3,5-triazine-2-carbonitrile,
 4-[(4-Chlorophenyl)amino]-6-pyrrolidin-1-yl-1,3,5-triazine-2-carbonitrile,
 5 4-[(4-Chlorophenyl)amino]-6-piperidin-1-yl-1,3,5-triazine-2-carbonitrile,
 4-[(4-Chlorophenyl)amino]-6-(ethylamino)-1,3,5-triazine-2-carbonitrile,
 4-[(4-Chlorophenyl)amino]-6-(3-hydroxypyrrolidin-1-yl)-1,3,5-triazine-2-carbonitrile,
 4-[(4-Chlorophenyl)amino]-6-[(2-piperidin-1-ylethyl)amino]-1,3,5-triazine-2-carbonitrile,
 4-[(4-Chlorophenyl)amino]-6-(4-phenylpiperidin-1-yl)-1,3,5-triazine-2-carbonitrile,
 10 4-[(3-Chlorobenzyl)amino]-6-(dimethylamino)-1,3,5-triazine-2-carbonitrile,
 4-Morpholin-4-yl-6-[(4-morpholin-4-ylphenyl)amino]-1,3,5-triazine-2-carbonitrile,
 4-(2,3-Dihydro-1,4-benzodioxin-6-ylamino)-6-morpholin-4-yl-1,3,5-triazine-2-
 carbonitrile,
 4-Morpholin-4-yl-6-(3-phenylpiperidin-1-yl)-1,3,5-triazine-2-carbonitrile,
 15 4-(1,4'-Bipiperidin-1'-yl)-6-morpholin-4-yl-1,3,5-triazine-2-carbonitrile,
 4-[4-(1H-Imidazol-1-yl)piperidin-1-yl]-6-morpholin-4-yl-1,3,5-triazine-2-carbonitrile,
 4-[4-(4-Chlorobenzoyl)piperidin-1-yl]-6-morpholin-4-yl-1,3,5-triazine-2-carbonitrile,
 4-[4-(5-Chloropyridin-2-yl)piperazin-1-yl]-6-morpholin-4-yl-1,3,5-triazine-2-carbonitrile,
 4-Morpholin-4-yl-6-{[3-(2-oxopyrrolidin-1-yl)propyl]amino}-1,3,5-triazine-2-carbonitrile,
 20 1-(4-Cyano-6-morpholin-4-yl-1,3,5-triazin-2-yl)-N,N-diethylpiperidine-3-carboxamide,
 4-[4-(2-Methoxyphenyl)piperazin-1-yl]-6-morpholin-4-yl-1,3,5-triazine-2-carbonitrile,
 N~2~-(4-Cyano-6-morpholin-4-yl-1,3,5-triazin-2-yl)-N~1~,N~1~-bis{4-[N-(4-cyano-6-
 morpholin-4-yl-1,3,5-triazin-2-yl)-N-isobutylglycyl]morpholin-3-yl}-N~2~-
 isobutylglycinamide,
 25 4-Morpholin-4-yl-6-[(2-pyridin-3-ylethyl)amino]-1,3,5-triazine-2-carbonitrile,
 4-[[2-(2-Furyl)ethyl]amino]-6-morpholin-4-yl-1,3,5-triazine-2-carbonitrile,
 4-[(4-Chlorophenyl)amino]-6-(4-methylpiperazin-1-yl)-1,3,5-triazine-2-carbonitrile,
 4-Azetidin-1-yl-6-[(4-chlorophenyl)amino]-1,3,5-triazine-2-carbonitrile,
 4-[(4-Chlorophenyl)amino]-6-morpholin-4-ylpyrimidine-2-carbonitrile,
 30 4-[(4-Methylcyclohexyl)amino]-6-morpholin-4-ylpyrimidine-2-carbonitrile,
 4-(4-Chlorophenoxy)-6-morpholin-4-ylpyrimidine-2-carbonitrile,
 4-[(4-Chlorophenyl)amino]-6-(dimethylamino)pyrimidine-2-carbonitrile,
 4-[(1-Methylpiperidin-4-yl)amino]-6-morpholin-4-ylpyrimidine-2-carbonitrile,
 4-(Cyclohexylamino)-6-morpholin-4-ylpyrimidine-2-carbonitrile,
 35 4-[(4-Chlorophenyl)amino]-6-pyrrolidin-1-ylpyrimidine-2-carbonitrile,
 4-[(6-Chloropyridin-3-yl)amino]-6-morpholin-4-ylpyrimidine-2-carbonitrile,

- 1-{6-[(4-Chlorophenyl)amino]-2-cyanopyrimidin-4-yl}-L-prolinamide,
4-(4-Aminopiperidin-1-yl)-6-[(4-chlorophenyl)amino]pyrimidine-2-carbonitrile,
4-[(4-Chlorophenyl)amino]-6-(4-pyrrolidin-1-yl)piperidin-1-ylpyrimidine-2-carbonitrile,
4-[(4-Chlorophenyl)amino]-6-[(3-pyrrolidin-1-ylpropyl)amino]pyrimidine-2-carbonitrile,
5 tert-Butyl 4-{6-[(4-chlorophenyl)amino]-2-cyanopyrimidin-4-yl}piperazine-1-carboxylate,
4-[(4-Chlorophenyl)amino]-6-(cyclopropylamino)pyrimidine-2-carbonitrile,
4-[(4-Chlorophenyl)amino]-6-piperazin-1-ylpyrimidine-2-carbonitrile,
(2S)-N-2-~{6-[(4-Chlorophenyl)amino]-2-cyanopyrimidin-4-yl}-N-1~,N-1~-bis[4-(N-
{6-[(4-chlorophenyl)amino]-2-cyanopyrimidin-4-yl}-L-leucyl)morpholin-3-yl]-L-
10 leucinamide,
5-Chloro-4-[(4-chlorophenyl)amino]-6-morpholin-4-ylpyrimidine-2-carbonitrile,
4-[(4-Chlorophenyl)amino]-5-methoxy-6-piperazin-1-ylpyrimidine-2-carbonitrile,
4-[(4-Chlorophenyl)amino]-5-methoxy-6-morpholin-4-ylpyrimidine-2-carbonitrile,
4-[(3S)-3-Aminopyrrolidin-1-yl]-6-[(4-chlorophenyl)amino]-5-methoxypyrimidine-2-
15 carbonitrile,
4-[(4-Chlorophenyl)amino]-6-{4-[3-(dimethylamino)propyl]piperazin-1-yl}-5-
methoxypyrimidine-2-carbonitrile,
4-[(4-Chlorophenyl)amino]-6-(dimethylamino)-5-methoxypyrimidine-2-carbonitrile,
4-[(4-Chlorophenyl)amino]-5-methoxy-6-(3-oxopiperazin-1-yl)pyrimidine-2-carbonitrile,
20 1-{6-[(4-Chlorophenyl)amino]-2-cyano-5-methoxypyrimidin-4-yl}piperidine-3-
carboxamide,
4-(4-Aminopiperidin-1-yl)-6-[(4-chlorophenyl)amino]-5-methoxypyrimidine-2-
carbonitrile,
5-Amino-4-[(4-chlorophenyl)amino]-6-morpholin-4-ylpyrimidine-2-carbonitrile,
25 5-Amino-4-[(4-Chlorophenyl)amino]-6-(ethylamino)pyrimidine-2-carbonitrile,
and pharmaceutically acceptable salts thereof.

7. A compound of formula (I) as defined in any one of claims 1 to 6 for use in therapy.
- 30 8. A pharmaceutical composition which comprises a compound of the formula (I) as
defined in any one of claims 1 to 6 or a pharmaceutically acceptable salt thereof and a
pharmaceutically acceptable diluent or carrier.
9. A method for producing inhibition of a cysteine protease in a mammal, such as man,
35 in need of such treatment, which comprises administering to said mammal an effective

amount of a compound of the present invention as defined in any one of claims 1 to 6, or a pharmaceutically acceptable salt thereof.

5

6
1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209
2210
2211
2212
2213
2214
2215
2216
221

ABSTRACT

The present invention relates to compounds and compositions for treating diseases associated with cysteine protease activity. The compounds are reversible inhibitors of cysteine proteases S, K, F, L and B. Of particular interest are diseases associated with Cathepsin S. In addition this invention also discloses processes for the preparation of such inhibitors.